INTERNATIONAL BACCALAUREATE

Mathematics: applications and interpretation

MAI

EXERCISES [MAI 3.14]

CROSS PRODUCT

Compiled by Christos Nikolaidis

A. Paper 1 questions (SHORT)

1. [Maximum mark: 8]

Let $\boldsymbol{a} = \begin{pmatrix} 2 \\ 0 \\ 3 \end{pmatrix}$ and $\boldsymbol{b} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$

(a) Find $\boldsymbol{a} \cdot \boldsymbol{b}$ and $\boldsymbol{b} \cdot \boldsymbol{a}$. [2]

(b) Find $\boldsymbol{a} \times \boldsymbol{b}$ and $\boldsymbol{b} \times \boldsymbol{a}$. [3]

(c) Show that $\boldsymbol{a} \times \boldsymbol{b}$ is perpendicular to both $\boldsymbol{a}$ and $\boldsymbol{b}$. [3]

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

2. [Maximum mark: 10]

Let $\boldsymbol{a} = \begin{pmatrix} 2 \\ 0 \\ 3 \end{pmatrix}$, $\boldsymbol{b} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ and $\boldsymbol{c} = \begin{pmatrix} 4 \\ 1 \\ 0 \end{pmatrix}$.

(a) Find $(\boldsymbol{a} \cdot \boldsymbol{b})\boldsymbol{c}$. [2]

(b) Find $(\boldsymbol{a} \times \boldsymbol{b}) \cdot \boldsymbol{c}$ and $\boldsymbol{a} \cdot (\boldsymbol{b} \times \boldsymbol{c})$. [6]

(c) Find $(\boldsymbol{a} \times \boldsymbol{b}) \times \boldsymbol{c}$. [2]

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

3. [Maximum mark: 8]

Consider the points $A(1,2,3)$, $B(3,0,5)$, $C(4,3,5))$.

(a) Find $\overrightarrow{AB} \times \overrightarrow{AC}$ [4]

(b) Find the area of the triangle ABC. [2]

The point $D(5,7,3)$ does not lie in the same plane with the triangle ABC.

The volume of the tetrahedron $ABCD$ is given by $\mathrm{V} = \frac{1}{6}\left|\overrightarrow{AD} \cdot (\overrightarrow{AB} \times \overrightarrow{AC})\right|$.

(c) Find the volume V. [2]

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

4. [Maximum mark: 5]

The position vectors of points P and Q are: $\boldsymbol{p} = 3\boldsymbol{i} + 2\boldsymbol{j} + \boldsymbol{k}$ and $\boldsymbol{q} = \boldsymbol{i} + 3\boldsymbol{j} - 2\boldsymbol{k}$

(a) Find the vector product $\boldsymbol{p} \times \boldsymbol{q}$. [3]

(b) Using your answer to part (a), or otherwise, find the area of the parallelogram with two sides $\overrightarrow{OP}$ and $\overrightarrow{OQ}$. [2]

....

....

....

....

....

....

....

....

....

....

5. [Maximum mark: 5]

(a) Find a vector perpendicular to the two vectors: $\overrightarrow{OP} = \vec{i} - 3\vec{j} + 2\vec{k}$, $\overrightarrow{OQ} = -2\vec{i} + \vec{j} - \vec{k}$. [3]

(b) If $\overrightarrow{OP}$ and $\overrightarrow{OQ}$ are position vectors for the points P and Q, use your answer to part (a), or otherwise, to find the area of the triangle OPQ. [2]

....

....

....

....

....

....

....

....

....

....

6. [Maximum mark: 5]

Let $\boldsymbol{a} = \begin{pmatrix} 2 \\ 1 \\ 0 \end{pmatrix}$, $\boldsymbol{b} = \begin{pmatrix} -1 \\ p \\ 6 \end{pmatrix}$ and $\boldsymbol{c} = \begin{pmatrix} 2 \\ -4 \\ 3 \end{pmatrix}$.

(a) Find $\boldsymbol{a} \times \boldsymbol{b}$. [3]

(b) Find the value of p, given that $\boldsymbol{a} \times \boldsymbol{b}$ is parallel to $\boldsymbol{c}$. [2]

....

....

....

....

....

....

....

....

....

....

....

7. [Maximum mark: 5]

Given that $\boldsymbol{a} = \boldsymbol{i} + 2\boldsymbol{j} - \boldsymbol{k}$, $\boldsymbol{b} = -3\boldsymbol{i} + 2\boldsymbol{j} + 2\boldsymbol{k}$ and $\boldsymbol{c} = 2\boldsymbol{i} - 3\boldsymbol{j} + 4\boldsymbol{k}$, find $(\boldsymbol{a} \times \boldsymbol{b}) \cdot \boldsymbol{c}$.

....

....

....

....

....

....

....

....

....

....

....

8. [Maximum mark: 5]

For the vectors $\boldsymbol{a} = 2\boldsymbol{i} + \boldsymbol{j} - 2\boldsymbol{k}$, $\boldsymbol{b} = 2\boldsymbol{i} - \boldsymbol{j} - \boldsymbol{k}$ and $\boldsymbol{c} = \boldsymbol{i} + 2\boldsymbol{j} + 2\boldsymbol{k}$, show that:

(a) $\boldsymbol{a} \times \boldsymbol{b} = -3\boldsymbol{i} - 2\boldsymbol{j} - 4\boldsymbol{k}$

(b) $(\boldsymbol{a} \times \boldsymbol{b}) \times \boldsymbol{c} = -(\boldsymbol{b} \bullet \boldsymbol{c})\boldsymbol{a}$

9. [Maximum mark: 4]

Find a vector that is perpendicular to the plane containing the lines L_1, and L_2, with vector equations

$L_1 : \boldsymbol{r} = \boldsymbol{i} + \boldsymbol{k} + \lambda\,(2\boldsymbol{i} + \boldsymbol{j} - 2\boldsymbol{k})$ $\quad$ $L_2 : \boldsymbol{r} = 3\boldsymbol{i} + 2\boldsymbol{j} + 2\boldsymbol{k} + \mu\,(\boldsymbol{j} + 3\boldsymbol{k})$

10. [Maximum mark: 6]

The parallelogram ABCD has vertices A(3,2,0), B(7,-1,-1), C(10,-3,0) and D(6,0,1). Calculate the area of the parallelogram.

....

....

....

....

....

....

....

....

....

....

....

11. [Maximum mark: 6]

Consider the points A(1, 2, –4), B(1, 5, 0) and C(6, 5, –12). Find the area of the triangle ABC.

....

....

....

....

....

....

....

....

....

....

....

....

....

12. [Maximum mark: 6]

Given that $\boldsymbol{a} = 2\boldsymbol{i} - \boldsymbol{j} - \boldsymbol{k}$, $\boldsymbol{b} = 2\boldsymbol{i} + \boldsymbol{j} - 2\boldsymbol{k}$ and $\boldsymbol{c} = -\boldsymbol{i} + \boldsymbol{j} - \boldsymbol{k}$ are the position vectors of the points A, B and C respectively, calculate the area of triangle ABC.

13. [Maximum mark: 6]

Given any two non-zero vectors $\boldsymbol{a}$ and $\boldsymbol{b}$, show that $|\boldsymbol{a} \times \boldsymbol{b}|^2 = |\boldsymbol{a}|^2 |\boldsymbol{b}|^2 - (\boldsymbol{a} \bullet \boldsymbol{b})^2$.

14. [Maximum mark: 7]

Three points A, B and C have coordinates (2, 1, –2), (2, –1, –1) and (1, 2, 2) respectively.

The vectors $\overrightarrow{OA}$, $\overrightarrow{OB}$ and $\overrightarrow{OC}$, where O is the origin, form three concurrent edges of a parallelepiped OAPBCQSR as shown in the following diagram.



(a) Find the coordinates of P, Q, R and S. [4]

(b) Calculate the volume, V, of the parallelepiped given that $V = |\overrightarrow{OA} \times \overrightarrow{OB} \bullet \overrightarrow{OC}|$ [3]

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

B. Paper 2 questions (LONG)

15. [Maximum mark: 13]

Three vectors $\boldsymbol{a}$, $\boldsymbol{b}$, and $\boldsymbol{c}$ (which are not coplanar) define a parallelepiped (or cuboid) and a tetrahedron (or triangular pyramid) as shown in the diagrams below.





It is given that

Volume of parallelepiped $= |(\boldsymbol{a} \times \boldsymbol{b}) \cdot \boldsymbol{c}|$

Volume of tetrahedron $= \dfrac{1}{6}|(\boldsymbol{a} \times \boldsymbol{b}) \cdot \boldsymbol{c}|$

Let $\boldsymbol{a} = \begin{pmatrix} 2 \\ 0 \\ 3 \end{pmatrix}$, $\boldsymbol{b} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$ and $\boldsymbol{c} = \begin{pmatrix} 4 \\ 1 \\ 0 \end{pmatrix}$.

(a) Find the area of the parallelogram and the area of triangle defined by $\boldsymbol{a}$ and $\boldsymbol{b}$. [6]

(b) Find the so-called **triple product** $(\boldsymbol{a} \times \boldsymbol{b}) \cdot \boldsymbol{c}$ [2]

(c) **Hence** find

(i) the volume of the parallelepiped defined by $\boldsymbol{a}$, $\boldsymbol{b}$, and $\boldsymbol{c}$.

(ii) the volume of the tetrahedron defined by $\boldsymbol{a}$, $\boldsymbol{b}$, and $\boldsymbol{c}$.

(iii) the height of the tetrahedron (or of the parallelepiped) corresponding to the base defined by $\boldsymbol{a}$ and $\boldsymbol{b}$ [5]

....

....

....

....

....

....

....

....

....

16. [Maximum mark: 12]

The coordinates of the points P, Q, R and S are $(4,1,-1)$, $(3,3,5)$, $(1,0,2c)$ and $(1,1,2)$. respectively. The vectors $\overrightarrow{QR}$ and $\overrightarrow{PR}$ are orthogonal (perpendicular).

(a) Find the value of c. [5]

(b) Evaluate $\overrightarrow{PS} \times \overrightarrow{PR}$. [4]

(c) Find the vector equation of the line *l* which passes through the point Q and is parallel to the vector PR. [3]

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

17. [Maximum mark: 10]

Consider the points $\mathrm{A}(1, 2, 1)$, $\mathrm{B}(0, -1, 2)$, $\mathrm{C}(1, 0, 2)$ and $\mathrm{D}(2, -1, -6)$.

(a) Find the vectors $\overrightarrow{\mathrm{AB}}$ and $\overrightarrow{\mathrm{BC}}$. [2]

(b) Calculate $\overrightarrow{\mathrm{AB}} \times \overrightarrow{\mathrm{BC}}$. [3]

(c) Hence, or otherwise find the area of triangle ABC. [2]

(d) Find a set of parametric equations for the line *L* through the point D and perpendicular to the triangle ABC. [3]

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

18. [Maximum mark: 11]

The points A, B, C, D have the following coordinates

A : (1, 3, 1) B : (1, 2, 4) C : (2, 3, 6) D : (5, – 2, 1).

(a) (i) Evaluate the vector product $\overrightarrow{AB} \times \overrightarrow{AC}$, giving your answer in terms of the unit vectors $\boldsymbol{i}, \boldsymbol{j}, \boldsymbol{k}$.

(ii) Find the area of the triangle ABC. [6]

(b) Find the vector equations of the lines

(i) L_1 which passes through D and is parallel to $\overrightarrow{AB}$.

(ii) L_2 which passes through D and A.

(iii) L_3 which passes through D and is perpendicular to the triangle ABC. [5]

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....

....