

INTERNATIONAL BACCALAUREATE
Mathematics: applications and interpretation

MAI

EXERCISES [MAI 4.9]
DISCRETE DISTRIBUTIONS
Compiled by Christos Nikolaidis

A. Paper 1 questions (SHORT)

1. [Maximum mark: 6]

The probability distribution of the discrete random variable X is given by the table

x	1	2	3	4	5
$P(X=x)$	0.4	0.2	0.15	0.15	0.1

Find

- (a) the expected value $E(X)$ of X . [2]
- (b) the mode of X . [1]
- (c) the median of X . [1]
- (d) the lower quartile Q_1 and the upper quartile Q_3 . [2]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

2. [Maximum mark: 6]

The probability distribution of the discrete random variable X is given by the table

x	1	2	3
$P(X=x)$	a	$2a$	b

Find the values of a and b given that $E(X) = 2.2$

.....

.....

.....

.....

.....

.....

.....

.....

3. [Maximum mark: 6]

The probability distribution of the discrete random variable X is given by the table

x	1	2	3
$P(X=x)$	0.2	0.4	0.4

Nikos selects a number at random.

If he selects 1 he earns 10 €. If he selects 2 he earns 5 €. If he selects 3 he loses 4 €

(a) Find the expected value of X . [3]

(b) Find the expected value of the profit for Nikos. Is the game fair? [3]

.....

.....

.....

.....

.....

.....

.....

.....

6. [Maximum mark: 6]

The probability distribution of the discrete random variable X is given by the following table.

x	1	2	3	4	5
$P(X=x)$	0.4	p	0.2	0.07	0.02

- (a) Find the value of p . [3]
 (b) Calculate the expected value of X . [3]

.....

7. [Maximum mark: 6]

A discrete random variable X has a probability distribution as shown in the table below.

x	0	1	2	3
$P(X=x)$	0.1	a	0.3	b

- (a) Find the value of $a + b$. [2]
 (b) Given that $E(X) = 1.5$, find the value of a and of b . [4]

.....

10. [Maximum mark: 6]

The following table shows the probability distribution of a discrete random variable X .

x	-1	0	2	3
$P(X=x)$	0.2	$10k^2$	0.4	$3k$

- (a) Find the value of k . [3]
 (b) Find the expected value of X . [3]

.....

11. [Maximum mark: 6]

A discrete random variable X has its probability distribution given by

$$P(X=x) = k(x+1), \text{ where } x \text{ is } 0, 1, 2, 3, 4.$$

- (a) Complete the following table showing the probability distribution for X
 (in terms of k) [2]

x	0	1	2	3	4
$P(X=x)$					

- (b) Show that $k = \frac{1}{15}$. [1]
 (c) Find $E(X)$. [3]

.....

12. [Maximum mark: 6]

The probability distribution of a discrete random variable X is defined by

$$P(X = x) = cx(5 - x), x = 1, 2, 3, 4.$$

(a) Find the value of c .

[3]

(b) Find $E(X)$.

[3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

13. [Maximum mark: 5]

The probability distribution of a discrete random variable X is given by

$$P(X = x) = k \left(\frac{2}{3}\right)^x, \text{ for } x = 0, 1, 2, \dots$$

Find the value of k .

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

15. [Maximum mark: 8]

A biased die with four faces is used in a game. A player pays 10 counters to roll the die. The table below shows the possible scores on the die, the probability of each score and the number of counters the player receives in return for each score.

Score	1	2	3	4
Probability	$\frac{1}{2}$	$\frac{1}{5}$	$\frac{1}{5}$	$\frac{1}{10}$
Number of counters player receives	4	5	15	n

(a) The player throws the die twice. Find the probability that

- (i) he has a total score of 3. (ii) he has a total score of 4.

[4]

(b) Find the value of n in order for the player to get an expected return of 9 counters per roll.

[4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

B. Paper 2 questions (LONG)

16. [Maximum mark: 15]

Two fair **four**-sided dice, one red and one green, are thrown. For each die, the faces are labelled 1, 2, 3, 4. The score for each die is the number which lands face down.

The sample space is shown below:

	1	2	3	4
1	•	•	•	•
2	•	•	•	•
3	•	•	•	•
4	•	•	•	•

(a) Write down the probability that two scores of 4 are obtained. [1]

Let X be the number of 4s that land face down.

(b) Complete the following probability distribution for X . [3]

x	0	1	2
$P(X = x)$			

(c) Find $E(X)$. [3]

Chris plays a game where he rolls the dice above.

If two 4s are obtained he wins 20€.

If only one 4 is obtained he wins 5€.

If no 4 is obtained he loses 2€

(d) Find the expected amount earned in one game. [3]

(e) If Chris plays this game 100 times find the amount he is expected to win. [2]

(f) If Chris plays this game twice find the probability that he earns 18€. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

17. [Maximum mark: 19]

Bag A contains 2 red balls and 3 green balls. Two balls are chosen at random from the bag without replacement. Let X denote the number of red balls chosen. The following table shows the probability distribution for X

X	0	1	2
$P(X=x)$	$\frac{3}{10}$	$\frac{6}{10}$	$\frac{1}{10}$

(a) Calculate $E(X)$, the mean number of red balls chosen. [3]

Bag B contains 4 red balls and 2 green balls. Two balls are chosen at random from bag B.

(b) (i) Draw a tree diagram to represent the above information, including the probability of each event.
 (ii) Hence find the probability distribution for Y , where Y is the number of red balls chosen. [8]

A standard die with six faces is rolled. If a 1 or 6 is obtained, two balls are chosen from bag A, otherwise two balls are chosen from bag B.

(c) Calculate the probability that two red balls are chosen. [5]
 (d) Given that two red balls are obtained, find the conditional probability that a 1 or 6 was rolled on the die. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

18. [Maximum mark: 16]

A **four-sided** die has three blue faces and one red face. The die is rolled.

Let B be the event a blue face lands down, and R be the event a red face lands down.


(a) Write down the values of

(i) $P(B)$

(ii) $P(R)$

[2]

(b) If the blue face lands down, the die is not rolled again. If the red face lands down, the die is rolled once again. This is represented by the following tree diagram, where p, s, t are probabilities.



Find the value of p , of s and of t .

[2]

Guisseppi plays a game where he rolls the die. If a blue face lands down, he scores 2 and is finished. If the red face lands down, he scores 1 and rolls one more time. Let X be the total score obtained.

(c) (i) Show that $P(X=3) = \frac{3}{16}$.

(ii) Find $P(X=2)$.

[3]

(d) (i) Construct a probability distribution table for X .

(ii) Calculate the expected value of X .

[5]

(e) If the total score is 3, Guisseppi wins \$10. If the total score is 2, Guisseppi gets nothing. He plays the game twice. Find the probability that he wins exactly \$10.

[4]

.....

.....

.....

.....

.....

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

19. [Maximum mark: 10]

John removes the labels from three cans of tomato soup and two cans of chicken soup in order to enter a competition, and puts the cans away. He then discovers that the cans are identical, so that he cannot distinguish between cans of tomato soup and chicken soup. Some weeks later he decides to have a can of chicken soup for lunch. He opens the cans at random until he opens a can of chicken soup. Let Y denote the number of cans he opens. Find

- (a) the possible values of Y , [1]
- (b) the probability of each of these values of Y , [4]
- (c) the expected value of Y . [2]
- (d) Write down the probability distribution of X if X denotes the number of cans he opens until he opens a can of tomato soup. [3]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....